Prognostic value of pre-transplant mean pulmonary arterial pressure in lung transplant recipients: a single-institution experience

Chi Young Kim¹, Ji Eun Park¹, Ah Young Leem¹, Joo Han Song¹, Song Yee Kim¹, Kyung Soo Chung¹, Eun Young Kim¹, Ji Ye Jung¹, Young Ae Kang¹, Young Sam Kim¹, Joon Chang¹, Jin Gu Lee², Hyo Chae Paik², Moo Suk Park¹

¹Division of Pulmonology, Department of Internal Medicine, Institute of Chest Diseases, Severance Hospital, Yonsei University College of Medicine, Seoul, Republic of Korea; ²Department of Thoracic & Cardiovascular Surgery, Severance Hospital, Yonsei University College of Medicine, Seoul, Republic of Korea

Contributions: (I) Conception and design: CY Kim, MS Park; (II) Administrative support: CY Kim, JY Park, SY Kim, MS Park; (III) Provision of study materials or patients: CY Kim, JY Park, SY Kim, MS Park; (IV) Collection and assembly of data: CY Kim, JY Park, SY Kim, MS Park; (V) Data analysis and interpretation: All authors; (VI) Manuscript writing: All authors; (VII) Final approval of manuscript: All authors.

Correspondence to: Moo Suk Park, MD, PhD. Division of Pulmonology, Department of Internal Medicine, Severance Hospital, Institute of Chest Diseases, Yonsei University College of Medicine, 50-1, Yonsei-ro, Seodaemun-gu, Seoul 120-752, Republic of Korea. Email: pms70@yuhs.ac.

Background: Currently, lung transplantation (LTX) is considered to be a curative treatment option in patients with end-stage lung disease. Although pulmonary hypertension (PH), confirmed by cardiac catheterization, is a prognostic factor in patients undergoing LTX, the prognostic value of PH in Asian lung transplant recipients remains uncertain. In this study, we aimed to determine whether PH before LTX may serve as a prognostic factor for survival in Asian patients.

Methods: The medical records of 50 patients [male, 27; female, 23; mean age, 51.0 (41.0–60.0) years], who received preoperative right heart catheterization (RHC) and echocardiography before single or double LTX at Severance Hospital between January 2010 and December 2014, were reviewed. The relationship between 1-year survival after LTX and PH [mean pulmonary arterial pressure (mPAP) ≥ 25 mmHg at rest] was evaluated.

Results: The mean right ventricular systolic pressure and mPAP were 48.5 (22.8) and 30.0 (24.0–40.0) mmHg. Of the 50 patients, 17 (34.0%) died within a year after LTX. The 1-year survival rate among patients with mPAP ≥ 25 mmHg (58.8%) was lower than the survival rate among patients with mPAP < 25 mmHg (87.5%). Pre-transplantation mPAP of ≥ 25 mmHg was associated with post-transplantation survival [hazard ratio (HR), 4.832; 95% confidence interval (CI), 1.080–21.608, $P=0.039$]. The presence of preoperative PH was also associated with an increased risk of postoperative complications.

Conclusions: Confirmation of PH via preoperative cardiac catheterization was associated with the prognosis of the patient after LTX. Clinicians should consider the necessity for early transplantation surgery before the mPAP reaches ≥ 25 mmHg.

Keywords: Lung transplantation (LTX); prognosis; hypertension; pulmonary

Submitted Sep 13, 2017. Accepted for publication Feb 12, 2018.

doi: 10.21037/jtd.2018.03.46

View this article at: <http://dx.doi.org/10.21037/jtd.2018.03.46>

Introduction

Management approaches for various intractable lung diseases have been transformed with the advancement of lung transplantation (LTX), which is one of the treatment options utilized at the endpoint of several lung diseases. Transplantation of other solid organs has produced better results and undergone greater improvement than LTX; thus, the outcomes of LTX remain among the worst overall for organ transplantation (1,2). However, the frequency and success rate of LTX have improved in recent years, with the increased success attributable to advances in post-LTX immunosuppression treatment (2,3); nevertheless, progress in LTX outcomes remains a challenging area of medicine. The survival rate associated with LTX has been increasing, such that the majority of deaths after LTX occur within the first year (4). Therefore, the identification of prognostic factors of 1-year mortality in lung transplant recipients might facilitate improvement of long term clinical outcomes after LTX.

There is a growing body of evidence that pulmonary hypertension (PH) associated with lung disease is a predictor of disease severity and mortality in patients awaiting LTX (5-7). Patients with PH suffer hypoxia, hypercapnia and respiratory acidosis, as a result of increased peripheral vascular resistance and right heart failure (8,9); these conditions can eventually lead to death. Previous studies have reported that PH, when confirmed by cardiac catheterization, is an independent predictive factor for survival after LTX (7,10-13). In addition, several studies have also analyzed post-LTX prognosis via echocardiography (14-16). However, almost all studies were exclusively designed for idiopathic pulmonary fibrosis (IPF) or other individual lung diseases. Moreover, there is a lack of consensus regarding predictive factors between IPF patients and patients who have other lung pathologies (17).

Despite the increasing popularity of LTX, there continue to be limited reports on the prognostic value of PH in lung transplant recipients. In addition, because of the small number of recipients of LTX in Asia, there are no reports in an Asian population regarding the clinical significance of PH in post-LTX survival. To determine the impact of PH on post-LTX survival, this study analyzed the characteristics and prognoses of lung transplant recipients with or without preoperative PH via echocardiography and right heart catheterization (RHC) in a Korean population.

Methods

Study design and population

We reviewed the medical records of 83 patients who underwent single or double LTX between January 2010 and July 2014 at Severance Hospital, Yonsei University, Republic of Korea. We excluded patients who did not undergo preoperative echocardiography or cardiac catheterization within 1 year prior to LTX, as well as those who missed follow-up echocardiography within 3 months after LTX. Thus, of the 83 patients whose records were initially reviewed, 50 were included in the study. From the database, we extracted all variables related to patient demographic characteristics and pre- and postoperative outcomes, in order to evaluate the relationship between PH and 1-year survival after LTX. Patient survival status was determined on the basis of hospital records. Donor data were collected from the Korean Network of Organ Sharing (KONOS) database. Additional variables included demographic characteristics, ischemic time, and donor-recipient size mismatch. All donated lungs from declared brain-dead patients were preserved in low-potassium dextran solution (Perfadex; Duraent Biologicals, Hyderabad, India). We divided the study subjects into two groups, according to their PH status.

RHC and echocardiography

RHC was performed at rest using a Swan-Ganz catheter. PH was defined as an increase in resting mean pulmonary arterial pressure (mPAP) of ≥ 25 mmHg, as assessed by RHC, using European Society of Cardiology and European Respiratory Society guidelines (18). Other echocardiographic variables were collected, such as ejection fraction (EF); right arterial pressure (RAP); regional wall motion abnormality (RWMA); ratio of mitral peak velocity of early filling (E) to early diastolic mitral annular velocity (E'), i.e., E/E'; and RV systolic pressure (RVSP). The characteristics of echocardiography were compared with RHC data, and correlation analysis was performed to investigate correlations with PH.

Statistical analysis

All data were analyzed using SPSS version 23 (IBM, Armonk, NY, USA). Data are expressed as mean values with standard deviations or median values with interquartile ranges (IQRs), as appropriate. Continuous

variables were compared by Student's *t*-test; categorical variables were compared by either the Chi-squared test or Fisher's exact test, as appropriate. Comparison of survival between patients with and without preoperative PH was performed by the log-rank test. The relationship between RHC and echocardiography-derived hemodynamics was assessed using the Pearson correlation. One-year survival following LTX was analyzed by Kaplan-Meier analysis; this measurement was compared between recipients with and without preoperative PH. Association of echocardiographic and hemodynamic parameters with clinical outcomes was evaluated by Cox proportional hazards regression analysis. Hazard ratios (HRs) and 95% confidence intervals (CIs) were calculated. In all statistical analyses, a two-tailed *P* value of <0.05 was considered to be statistically significant.

Ethical approval

The study protocol was approved by the institutional review board of Severance Hospital (approval No. 2013-0522-019). In addition, this study was conducted in compliance with the principles set in the 2000 Declaration of Helsinki, as well as the 2008 Declaration of Istanbul.

Results

Baseline characteristics of lung transplant recipients

Table 1 describes the baseline characteristics of all patients who were included in this analysis, comprising 50 [27 males (54%) and 23 females (46%)] patients. The mean age of the patients was 51.0 (41.0–60.0) years; 47 (94.0%) patients underwent double LTX. On the basis of mPAP values, 34 patients (68.0%) were classified into the PH group; the remaining patients comprised the non-PH group. Comparison of demographic, hemodynamic, and pulmonary function characteristics among lung transplant recipients was performed, according to PH status. The distribution of underlying lung diseases was as follows: IPF, 22 patients (44.0%); lymphangioleiomyomatosis, 8 patients (16.0%); connective tissue disease, 6 patients (12.0%); bronchiectasis, 5 patients (10.0%); bronchiolitis obliterans after stem cell transplantation, 4 patients (8.0%); diffuse panbronchiolitis, 2 patients (4.0%); and others [chronic obstructive pulmonary disease, other pulmonary fibrosis, and pulmonary Langerhans cell histiocytosis, 1 patient each (2.0%)]. There was no statistically significant difference in the distribution of the underlying diseases. Pulmonary

function tests were administered before LTX; lung transplant recipients exhibited underlying lung diseases with severe restrictive patterns, as evidenced by the relatively low mean values of functional vital capacity (45.6%±16.3%) and diffusing capacity of the lung for carbon monoxide (DL_{CO}; 37.8%±18.8%). The mean values of preoperative RVSP and mPAP (determined by echocardiography and RHC, respectively) were 48.5±22.8 and 30.0 (24.0–40.0) mmHg, respectively. Comparison of baseline characteristics revealed that there were significant differences between the two groups regarding most hemodynamic variables, suggesting that hemodynamic variables may act as a prognostic factor; however, the only significant prognostic factors in both univariate analysis and multivariate analysis were mPAP and RAP.

Clinical predictors of 1-year survival after LTX

Of the 50 patients in the study, 17 (34.0%) died within 1 year after LTX. The 1-year survival rate of lung transplant recipients was higher in the PH group; the survival rates of patients with and without PH were 58.8% and 87.0%, respectively (log-rank test, *P*=0.030; Figure 1). Table 2 presents the results of univariate analysis of predictive values for 1-year survival after LTX. Old age, male sex, higher BMI, RAP, and mPAP were identified as independent prognostic factors.

The results of multivariate Cox proportional hazards analysis of age, sex, BMI, underlying disease (IPF *vs.* non-IPF), and mPAP—all of which were identified as significant prognostic factors in univariate analysis—revealed an HR of 4.832 (95% CI, 1.080–21.608; *P*=0.039) for survival in patients with mPAP ≥25 mmHg. These results strongly suggest that preoperative PH significantly affects the chances of 1-year survival (Table 3).

Postoperative complications among lung transplant recipients

Table 4 presents the comparison of postoperative complications, occurring within 3 months of LTX, in lung transplant recipients. Complications such as sepsis, acute kidney injury, acute rejection, postoperative bleeding, and grade 3 primary graft dysfunction were more frequent in the PH group than in the non-PH group. The proportion of complications was higher in the PH group than in the non-PH group.

Table 1 Comparison of baseline characteristics of patients with and without pulmonary hypertension before lung transplantation (N=50)

Variables	Patient without PH (n=16)	Patients with PH (n=34)	Total (n=50)	P value
Age, years	51.5 (39.5–58.0)	51.0 (43.0–60.0)	51.0 (41.0–60.0)	0.755
Male gender, n (%)	7 (43.8)	20 (58.8)	27 (54.0)	0.488
Body mass index, kg/m ²	18.6 (3.7)	20.6 (4.0)	19.9 (4.0)	0.097
Smoking, pack years	22.0 (20.0–26.0)	40.0 (16.0–53.0)	22.0 (18.5–45.0)	0.420
Underlying diseases, n (%)				0.133
IPF, n (%)	10 (62.5)	12 (35.3)	22 (44.0)	
Non-IPF, n (%) [*]	6 (37.5) [†]	22 (64.7) [‡]	28 (56.0)	
Double LTX, n (%)	16 (100.0)	31 (91.2)	47 (94.0)	0.557
Cr, mg/dL	0.6 (0.2)	0.6 (0.2)	0.6 (0.2)	0.951
Albumin, mg/dL	3.6 (2.9–4.2)	3.8 (3.0–4.2)	3.8 (3.0–4.2)	0.884
NT-ProBNP, pg/mL	62.0 (23.0–167.5)	49.0 (23.5–895.0)	51.0 (23.0–714.0)	0.623
Coronary disease, n (%)	2 (12.5)	8 (23.5)	10 (20.0)	0.681
DM, n (%)	3 (18.8)	6 (17.6)	9 (18.0)	1.000
6MWT, M	290.0 (170.0–317.5)	204.1 (125.9)	222.7 (128.3)	0.151
FEV ₁ , % predicted	44.6 (17.6)	40.7 (20.4)	41.9 (19.4)	0.509
FVC, % predicted	43.2 (15.5)	46.8 (16.8)	45.6 (16.3)	0.483
DLCO, mL/mmHg/min	45.73 (20.9)	33.72 (16.5)	37.8 (18.8)	0.299
Pre-op MV, n (%)	2 (12.5)	5 (14.7)	7 (14.0)	1.000
Pre-op OP ECMO, n (%)	0 (0.0)	2 (5.9)	2 (4.0)	0.829
Echocardiography data				
EF, %	65.1(8.0)	63.7 (9.3)	64.1 (8.9)	0.618
RWMA, n (%)	1 (6.3)	2 (5.9)	3 (6.0)	1.000
RAP, mmHg	5.0 (1.3)	8.0 (4.9)	7.2 (4.3)	0.056
RVSP, mmHg	39.8 (14.1)	52.2 (24.9)	48.5 (22.8)	0.037
E/E' ratio	8.8 (2.0)	9.4 (2.8)	9.2 (2.5)	0.461
Hemodynamic data				
mPAP, mmHg	21.1 (19.0–23.5)	32.5 (28.0–40.6)	30.0 (24.0–40.0)	<0.001
RAP, mmHg	6.0 (3.5–8.5)	10.0 (6.0–15.0)	8.0 (5.0–13.0)	0.008
RVP, mmHg	29.5 (27.0–36.5)	43.0 (32.0–57.0)	36.5 (27.0–52.0)	0.010
PCWP, mmHg	11.0 (8.5–12.0)	12.0 (10.0–21.0)	11.0 (9.5–20.5)	0.052
CO, L/min	4.5 (1.3)	5.1 (1.1)	4.7 (1.2)	0.042
CI, L/min/m ²	2.8 (0.8)	3.4 (0.8)	3.0 (0.9)	0.092

Table 1 (continued)

Table 1 (continued)

Variables	Patient without PH (n=16)	Patients with PH (n=34)	Total (n=50)	P value
Donor				
Age, years	44.0 (34.5–51.0)	41.0 (37.0–47.0)	42.0 (37.0–50.0)	0.487
Male gender, n (%)	8 (50.0)	25 (73.5)	33 (66.0)	0.139
Body mass index, kg/m ²	22.4 (2.9)	22.4 (3.0)	22.4 (2.9)	0.982
Smoking, pack years	9.1 (2.5–13.8)	8.8 (3.2–15.0)	10.0 (2.5–15.0)	0.799
Ischemic time, min	219.5 (160.0–265.5)	215.0 (181.5–251.0)	216.0 (176.0–254.5)	0.875
Size mismatch, n (%)	4 (25.0)	7 (20.6)	11 (22.0)	1.000

Values are expressed as mean (SD) or median (IQR). *, the distribution of underlying lung diseases; †, bronchiolitis obliterans after stem cell transplantation, 4 (25.0%); lymphangioleiomyomatosis, 1 (6.3%); connective tissue disease, 1 (6.3%); ‡, lymphangioleiomyomatosis, 7 (20.6%); bronchiectasis, 5 (14.7%); connective tissue disease, 5 (14.7%); diffuse panbronchiolitis, 2 (5.9%); chronic obstructive pulmonary disease, 1 (2.9%); pulmonary Langerhans cell histiocytosis, 1 (2.9%); other pulmonary fibrosis, 1 (2.9%). IPF, idiopathic pulmonary fibrosis; Cr, serum creatinine; NT-proBNP, n-terminal pro-brain natriuretic peptide; Coronary disease, need a medical treatment in preoperative coronary angiography; DM, diabetes mellitus; 6MWT, 6 minute walk test; FEV₁, forced expiratory volume in 1 s; FVC, forced vital capacity; DLCO, diffusion capacity for carbon monoxide; pre-op, preoperative; MV, mechanical ventilation; ECMO, extracorporeal membrane oxygenation; EF, ejection fraction; RWMA, regional wall motion abnormality; RAP, right arterial pressure; RVSP, right ventricular systolic pressure; E/E', ratio of mitral peak velocity of early filling (E) to early diastolic mitral annular velocity (E'); mPAP, mean pulmonary arterial pressure; RVP, right ventricular pressure; PCWP, pulmonary capillary wedge pressure; CO, cardiac output; CI, cardiac index; SD, standard deviation; IQR, interquartile range.

Figure 1 Kaplan-Meier survival curve for 1-year mortality after lung transplantation, according to pulmonary hypertension status. PAP, pulmonary arterial pressure.

Discussion

This study investigated the association between 1-year mortality and preoperative PH among patients who underwent LTX, revealing higher rates of 1-year mortality and postoperative complications among patients with preoperative PH, relative to patients without preoperative PH. Thus, the presence of preoperative PH was associated with an increased risk of postoperative complications and mortality.

The presence of PH is a significant prognostic factor in patients with end-stage lung diseases. Among patients with COPD, up to 90% of those with stage IV disease (according to the Global Initiative for Chronic Obstructive Lung Disease guidelines) demonstrate mPAP of >20 mmHg (8). In addition, the prevalence of PH among patients with IPF ranges from 31.6% to 84% (10,13,19). Similarly, in various lung diseases, most patients develop PH with disease progression (17). It is interesting to note that, among lung transplant recipients in the present study, preoperative PH was associated with poor

Table 2 Results of univariate analysis of prognostic indicators for 1-year mortality after lung transplantation

Variables	HR (95% CI)	P value
Age, years	1.085 (1.030–1.143)	0.002
Male gender, n (%)	0.416 (0.146–1.181)	0.099
Body mass index, kg/m ²	1.249 (1.113–1.401)	<0.001
Smoking, pack years	1.018 (0.985–1.052)	0.292
Underlying disease		0.123
IPF	Reference range	
Non-IPF*	0.467 (0.177–1.229)	
Double LTX	2.913 (0.663–12.794)	0.157
Coronary disease	0.481 (0.159–1.368)	0.170
DM	1.444 (0.470–4.439)	0.521
6MWT, M	0.999 (0.995–1.004)	0.766
Echocardiography data		
EF, %	1.019 (0.963–1.079)	0.514
RWMA, N (%)	0.801 (0.106–6.047)	0.851
RAP, mmHg	1.044 (0.943–1.156)	0.409
RVSP, mmHg	1.005 (0.984–1.027)	0.615
E/E' ratio	1.116 (0.944–1.320)	0.200
Hemodynamic data		
mPAP ≥25, mmHg	4.444 (1.015–19.454)	0.048
RAP, mmHg	1.086 (1.021–1.155)	0.008
RVP, mmHg	1.008 (0.989–1.027)	0.407
PCWP, mmHg	1.035 (0.985–1.088)	0.177
CO, L/min	1.278 (0.717–2.277)	0.405
CI, L/min/m ²	0.659 (0.240–1.808)	0.659
Pre-op MV	2.491 (0.88–7.686)	0.112
Pre-op ECMO	1.923 (0.254–14.557)	0.527
Donor		
Age, years	0.983 (0.944–1.025)	0.425
Male gender	0.612 (0.197–1.898)	0.395
Body mass index, kg/m ²	0.948 (0.794–1.131)	0.551

Table 2 (continued)**Table 2** (continued)

Variables	HR (95% CI)	P value
Smoking, pack years	0.933 (0.825–1.054)	0.265
Ischemic time, min	1.004 (0.997–1.012)	0.216
Size mismatch	1.972 (0.727–5.347)	0.182

*, the distribution of underlying lung diseases: lymphangioleiomyomatosis, 8 (28.6%); connective tissue disease, 6 (21.4%); bronchiectasis, 5 (17.6); bronchiolitis obliterans after stem cell transplantation, 4 (14.3%); diffuse panbronchiolitis, 2 (7.1); chronic obstructive pulmonary disease, 1 (3.6%); pulmonary Langerhans cell histiocytosis, 1 (3.6%); other pulmonary fibrosis, 1 (3.6%). HR, hazard ratio; CI, confidence interval; mPAP, mean pulmonary arterial pressure; IPF, idiopathic pulmonary fibrosis; Coronary disease, need a medical treatment in preoperative coronary angiography; DM, diabetes mellitus; 6MWT, 6 minute walk test; EF, ejection fraction; RWMA, regional wall motion abnormality; RAP, right arterial pressure; RVSP, right ventricular systolic pressure; E/E', ratio of mitral peak velocity of early filling (E) to early diastolic mitral annular velocity (E'); RVP, right ventricular pressure; PCWP, pulmonary capillary wedge pressure; CO, cardiac output; pre-op, preoperative; MV, mechanical ventilation; ECMO, extracorporeal membrane oxygenation; PGD, primary graft dysfunction.

Table 3 Results of multivariate analysis of prognostic indicators for 1-year mortality after lung transplantation

Variables	HR (95% CI)	P value
Age, years	1.067 (1.006–1.132)	0.030
Male gender	1.007 (0.244–4.157)	0.992
BMI, kg/m ²	1.146 (1.003–1.309)	0.045
IPF vs. non-IPF	0.822 (0.190–3.566)	0.794
RAP, mmHg	1.028 (0.949–1.114)	0.492
mPAP ≥25, mmHg	4.832 (1.080–21.608)	0.039

HR, hazard ratio; CI, confidence interval; BMI, body mass index; IPF, idiopathic pulmonary fibrosis; RAP, right arterial pressure; mPAP, mean pulmonary arterial pressure.

Table 4 Postoperative complications within 3 months in lung transplant recipients

Variables	Patient without PH (n=16)	Patient with PH (n=34)	Total (n=50)	P value
Pneumonia, n (%)	12 (75.0)	28 (82.4)	40 (80.0)	0.960
Sepsis, n (%)	3 (18.8)	15 (44.1)	18 (36.0)	0.081
AKI, n (%)	6 (37.5)	17 (50.0)	23 (46.0)	0.307
RRT, n (%)	2 (12.5)	13 (38.2)	15 (30.0)	0.113
Post-op bleeding, n (%)	2 (12.5)	6 (17.6)	8 (16.0)	0.960
Acute rejection, n (%)	0 (0.0)	4 (11.8)	4 (8.0)	0.383
Total HOD, days	41.4 (8.0–97.0)	59.5(23.0–159.0)	48.9 (25.3–74.3)	0.139
ICU stay after operation, days	7.0 (6.0–22.5)	8.5 (7.0–29.0)	8.0 (6.0–26.0)	0.535
Post-op MV, days	4.0 (3.0–19.0)	8.0 (4.0–19.0)	8.0 (4.0–19.0)	0.549
PGD grade 3, n (%)	1 (6.3)	7 (20.6)	8 (16.0)	0.197

Values are expressed as mean (SD) or median (IQR). SD, standard deviation; IQR, interquartile range; PH, pulmonary hypertension; AKI, acute kidney injury; RRT, renal replacement therapy; post-op, post-operative; HOD, day of hospital stay; ICU, intensive care unit; MV, mechanical ventilation; PGD, primary graft dysfunction.

prognosis for 1-year survival post-LTX.

Several studies and guidelines have defined PH as mPAP ≥ 25 mmHg in patients with or without underlying diseases; in addition, RHC has been utilized as the “gold standard” for PH diagnosis (9,17,20). In the past, chest X-ray imaging and echocardiography were used to determine the presence and severity of PH (17). However, given its relatively high accuracy in the evaluation of PH status among lung transplant recipients, RHC is increasingly used in pre- and postoperative evaluation of PH. Although RHC is the most accurate method to diagnose PH, it is an invasive procedure. Therefore, echocardiography is preferred as the initial screening modality for diagnosis of PH. A few studies have used echocardiography to investigate mortality among lung transplant recipients with poor RV function (15,16,21).

However, this diagnostic technology requires estimates of tricuspid regurgitation jet velocity and is associated with underestimation of pressures in patients with high PAP, as well as overestimation of PAP in patients without significant elevation (22,23). In the present study, the presence of PH was found in 68% of patients by RHC, whereas it was found in 77% of patients by use of echocardiography [defined as a pulmonary artery systolic pressure (PASP) of >35 mmHg]. Unfortunately, there was no correlation between echocardiography and survival (HR 1.005; 95% CI, 0.984–1.027; $P=0.615$). There was a positive correlation was observed between PASP estimated by echocardiography and measured by RHC ($r=0.584$, $P<0.001$). Although

there is a good correlation in baseline PASP between echocardiography and RHC, echocardiographic PH measurements alone are not sufficient to monitor changes in PASP or progression of PH, as described in previous studies (24). Although it may serve as a good screening tool, echocardiographic PH measurement alone is not sufficient to predict mortality or postoperative outcome. Further large prospective cohort studies are required regarding the diagnosis of PH by echocardiography.

In contrast to the present findings, Whitson *et al.* reported a lack of strong evidence for a relationship between preoperative PH and postoperative mortality (17). Furthermore, several studies have been unable to identify a statistically significant difference in survival between patients with PH and patients without PH (25,26). Because of these inconsistent results, the impact of preoperative PH on postoperative mortality is still controversial. However, most previous studies, including a large prospective cohort study, have reported high mortality or poor outcomes among lung transplant recipients with PH-associated lung disease (7,10–13,17,19,27–30). In addition, the variation in the impact of PH on post-transplantation mortality might be attributable to the heterogeneity of end-stage lung diseases among the present study population. In a recent publication, Gries *et al.* also reported differences in patient survival related to the underlying lung disease (30).

A growing body of evidence suggests that age at operation, sex, BMI, preoperative albumin level, kidney

function, ischemic time, and size mismatch are significant prognostic factors for survival after LTX (7,21,30-32). The present findings differ from those of some previous reports regarding survival after LTX. In this study, significant prognostic factors for post-LTX survival were limited to age, sex, BMI, and PH (mPAP ≥ 25 mmHg). Upon multivariate analysis—performed to exclude the effect of non-PH factors, including underlying disease (IPF or non-IPF), on survival—we found that the severity of PH, as defined by mPAP, was a prognostic factor for 1-year survival in lung transplant recipients.

Unfortunately, most studies that have evaluated the risk factors of post-LTX mortality were performed in North America and Europe. As the medical environment and causes of LTX in Korea are very different relative to the previous studies, these differences may also affect the prognoses of patients undergoing LTX. Furthermore, the mortality rate in patients awaiting LTX in Korea was higher than in other countries (33-35). This may be attributable to the lack of donated lungs and the low utilization rate of donated lungs from brain-dead patients, which are the major obstacles to LTX in Korea, relative to other countries (36).

To our knowledge, this was the first study to evaluate the clinical significance of post-LTX PH in Asian patients. Our results clearly demonstrated a significant correlation between mPAP and 1-year survival after LTX. In addition, the rate and severity of postoperative complications, such as sepsis, multi-organ dysfunction, postoperative bleeding, postoperative gastrointestinal bleeding, and grade 3 primary graft dysfunction, were higher among patients with PH than among patients without PH. For these reasons, we believe that PH is a significant prognostic factor for post-LTX survival.

In chronic lung diseases associated with PH, RV function is strongly influenced by the afterload (i.e., the underlying lung disease status) (17). Additionally, PH also causes distortion of cardiac geometry; however, post-LTX echocardiographic findings have demonstrated reverse remodeling of the distorted cardiac geometry (37). Thus, for a variety of reasons, severe pulmonary disease affects both the right and left ventricles and leads to PH (38), which is considered to be associated with increased mortality. There is an increasing effort to produce evidence supporting this claim in specific lung diseases, such as cystic fibrosis (26,39,40). Collectively, these findings suggest that the presence of PH before LTX might affect the prognosis in lung transplant recipients. Therefore, further studies,

involving a large number of patients with various end-stage lung diseases, are required to evaluate the prognostic implication of preoperative PH in the clinical outcome of LTX.

Limitations of the present study include its relatively small sample size and retrospective study design solely involving medical record review. In addition, there is no apparent difference in the prognosis between severe PH and mild to moderate PH; this requires additional investigation. Despite these limitations, the strength of this study is that this was the first study to evaluate the prognostic significance of preoperative PH in Asian patients; moreover, the data were derived from a single center cohort. Considering that differences in ethnicity influence the outcome in patients with end-stage lung disease, data from our study could provide useful information regarding the prognosis of Asian patients undergoing LTX.

In conclusion, patients with PH exhibited higher 1-year mortality after LTX, relative to patients without PH. PH associated with lung disease has been shown to be a predictor of the severity of disease and mortality in patients awaiting LTX. Thus, the evaluation of PH by preoperative RHC might be essential in patients who are awaiting LTX. Larger prospective studies are required to verify the benefit of early transplantation in patients with PH, following evaluation by echocardiography and RHC. Furthermore, earlier transplantation should be considered before the occurrence of end-stage lung disease in patients with PH.

Acknowledgements

None.

Footnote

Conflicts of Interest: The authors have no conflicts of interest to declare.

Ethical Statement: The study protocol was approved by the institutional review board of Severance Hospital (approval No. 2013-0522-019). In addition, this study was conducted in compliance with the principles set in the 2000 Declaration of Helsinki, as well as the 2008 Declaration of Istanbul.

References

1. Christie JD, Edwards LB, Kucheryavaya AY, et al. The

- Registry of the International Society for Heart and Lung Transplantation: Twenty-eighth Adult Lung and Heart-Lung Transplant Report--2011. *J Heart Lung Transplant* 2011;30:1104-22.
2. Studer SM, Levy RD, McNeil K, et al. Lung transplant outcomes: a review of survival, graft function, physiology, health-related quality of life and cost-effectiveness. *Eur Respir J* 2004;24:674-85.
 3. Strueber M, Warnecke G, Fuge J, et al. Everolimus Versus Mycophenolate Mofetil De Novo After Lung Transplantation: A Prospective, Randomized, Open-Label Trial. *Am J Transplant* 2016;16:3171-80.
 4. Sabashnikov A, Weymann A, Mohite PN, et al. Risk factors predictive of one-year mortality after lung transplantation. *Eur J Cardiothorac Surg* 2014;46:e82-8.
 5. Hayes D Jr, Black SM, Tobias JD, et al. Prevalence of Pulmonary Hypertension and its Influence on Survival in Patients With Advanced Chronic Obstructive Pulmonary Disease Prior to Lung Transplantation. *Copd* 2016;13:50-6.
 6. Hayes D Jr, Black SM, Tobias JD, et al. Influence of pulmonary hypertension on survival in advanced lung disease. *Lung* 2015;193:213-21.
 7. Hayes D Jr, Black SM, Tobias JD, et al. Influence of Pulmonary Hypertension on Patients With Idiopathic Pulmonary Fibrosis Awaiting Lung Transplantation. *Ann Thorac Surg* 2016;101:246-52.
 8. Wrobel JP, Thompson BR, Williams TJ. Mechanisms of pulmonary hypertension in chronic obstructive pulmonary disease: a pathophysiologic review. *J Heart Lung Transplant* 2012;31:557-64.
 9. Seeger W, Adir Y, Barbera JA, et al. Pulmonary hypertension in chronic lung diseases. *J Am Coll Cardiol* 2013;62:D109-16.
 10. Shorr AF, Wainright JL, Cors CS, et al. Pulmonary hypertension in patients with pulmonary fibrosis awaiting lung transplant. *Eur Respir J* 2007;30:715-21.
 11. Hayes D Jr, Tumin D, Budev MM, et al. Adverse outcomes associated with pulmonary hypertension in chronic obstructive pulmonary disease after bilateral lung transplantation. *Respir Med* 2017;128:102-8.
 12. Andersen KH, Schultz HH, Nyholm B, et al. Pulmonary hypertension as a risk factor of mortality after lung transplantation. *Clin Transplant* 2016;30:357-64.
 13. Nadrous HF, Pellikka PA, Krowka MJ, et al. Pulmonary hypertension in patients with idiopathic pulmonary fibrosis. *Chest* 2005;128:2393-9.
 14. Rudski LG, Lai WW, Afilalo J, et al. Guidelines for the echocardiographic assessment of the right heart in adults: a report from the American Society of Echocardiography endorsed by the European Association of Echocardiography, a registered branch of the European Society of Cardiology, and the Canadian Society of Echocardiography. *J Am Soc Echocardiogr* 2010;23:685-713; quiz 86-8.
 15. Kusunose K, Tsutsui RS, Bhatt K, et al. Prognostic value of RV function before and after lung transplantation. *JACC Cardiovasc Imaging* 2014;7:1084-94.
 16. Porteous MK, Ky B, Kirkpatrick JN, et al. Diastolic Dysfunction Increases the Risk of Primary Graft Dysfunction after Lung Transplant. *Am J Respir Crit Care Med* 2016;193:1392-400.
 17. Hayes D Jr, Higgins RS, Black SM, et al. Effect of pulmonary hypertension on survival in patients with idiopathic pulmonary fibrosis after lung transplantation: an analysis of the United Network of Organ Sharing registry. *J Heart Lung Transplant* 2015;34:430-7.
 18. Galie N, Humbert M, Vachiery JL, et al. 2015 ESC/ERS Guidelines for the diagnosis and treatment of pulmonary hypertension: The Joint Task Force for the Diagnosis and Treatment of Pulmonary Hypertension of the European Society of Cardiology (ESC) and the European Respiratory Society (ERS): Endorsed by: Association for European Paediatric and Congenital Cardiology (AEPC), International Society for Heart and Lung Transplantation (ISHLT). *Eur Respir J* 2015;46:903-75.
 19. Lettieri CJ, Nathan SD, Barnett SD, et al. Prevalence and outcomes of pulmonary arterial hypertension in advanced idiopathic pulmonary fibrosis. *Chest* 2006;129:746-52.
 20. Rosenkranz S, Preston IR. Right heart catheterisation: best practice and pitfalls in pulmonary hypertension. *Eur Respir Rev* 2015;24:642-52.
 21. Armstrong HF, Schulze PC, Kato TS, et al. Right ventricular stroke work index as a negative predictor of mortality and initial hospital stay after lung transplantation. *J Heart Lung Transplant* 2013;32:603-8.
 22. Brecker SJ, Gibbs JS, Fox KM, et al. Comparison of Doppler derived haemodynamic variables and simultaneous high fidelity pressure measurements in severe pulmonary hypertension. *Br Heart J* 1994;72:384-9.
 23. Arcasoy SM, Christie JD, Ferrari VA, et al. Echocardiographic assessment of pulmonary hypertension in patients with advanced lung disease. *Am J Respir Crit Care Med* 2003;167:735-40.
 24. Opitz C, Rosenkranz S, Ghofrani HA, et al. ESC guidelines 2015 pulmonary hypertension: diagnosis and

- treatment. *Dtsch Med Wochenschr* 2016;141:1764-9.
25. Andersen KH, Iversen M, Kjaergaard J, et al. Prevalence, predictors, and survival in pulmonary hypertension related to end-stage chronic obstructive pulmonary disease. *J Heart Lung Transplant* 2012;31:373-80.
 26. Belle-van Meerkerk G, Cramer MJ, Kwakkel-van Erp JM, et al. Pulmonary hypertension is a mild comorbidity in end-stage cystic fibrosis patients. *J Heart Lung Transplant* 2013;32:609-14.
 27. Armstrong HF, Schulze PC, Bacchetta M, et al. Impact of pulmonary hypertension on exercise performance in patients with interstitial lung disease undergoing evaluation for lung transplantation. *Respirology* 2014;19:675-82.
 28. Kimura M, Taniguchi H, Kondoh Y, et al. Pulmonary hypertension as a prognostic indicator at the initial evaluation in idiopathic pulmonary fibrosis. *Respiration* 2013;85:456-63.
 29. Teramachi R, Taniguchi H, Kondoh Y, et al. Progression of mean pulmonary arterial pressure in idiopathic pulmonary fibrosis with mild to moderate restriction. *Respirology* 2017;22:986-90.
 30. Singh VK, Patricia George M, Gries CJ. Pulmonary hypertension is associated with increased post-lung transplant mortality risk in patients with chronic obstructive pulmonary disease. *J Heart Lung Transplant* 2015;34:424-9.
 31. Gries CJ, Bhadriraju S, Edelman JD, et al. Obese patients with idiopathic pulmonary fibrosis have a higher 90-day mortality risk with bilateral lung transplantation. *J Heart Lung Transplant* 2015;34:241-6.
 32. Chandrashekar S, Keller CA, Kremers WK, et al. Weight loss prior to lung transplantation is associated with improved survival. *J Heart Lung Transplant* 2015;34:651-7.
 33. Glanville AR, Estenne M. Indications, patient selection and timing of referral for lung transplantation. *Eur Respir J* 2003;22:845-52.
 34. Paik HC, Haam SJ, Lee DY, et al. The fate of patients on the waiting list for lung transplantation in Korea. *Transplant Proc* 2012;44:865-9.
 35. Orens JB, Estenne M, Arcasoy S, et al. International guidelines for the selection of lung transplant candidates: 2006 update--a consensus report from the Pulmonary Scientific Council of the International Society for Heart and Lung Transplantation. *J Heart Lung Transplant* 2006;25:745-55.
 36. Paik HC, Haam SJ, Lee DY, et al. Donor evaluation for lung transplantation in Korea. *Transplant Proc* 2012;44:870-4.
 37. Kasimir MT, Seebacher G, Jaksch P, et al. Reverse cardiac remodelling in patients with primary pulmonary hypertension after isolated lung transplantation. *Eur J Cardiothorac Surg* 2004;26:776-81.
 38. Magruder JT, Shah AS, Crawford TC, et al. Simulated Regionalization of Heart and Lung Transplantation in the United States. *Am J Transplant* 2017;17:485-95.
 39. Tonelli AR, Fernandez-Bussy S, Lodhi S, et al. Prevalence of pulmonary hypertension in end-stage cystic fibrosis and correlation with survival. *J Heart Lung Transplant* 2010;29:865-72.
 40. Scarsini R, Prioli MA, Milano EG, et al. Hemodynamic predictors of long term survival in end stage cystic fibrosis. *Int J Cardiol* 2016;202:221-5.

Cite this article as: Kim CY, Park JE, Leem AY, Song JH, Kim SY, Chung KS, Kim EY, Jung JY, Kang YA, Kim YS, Chang J, Lee JG, Paik HC, Park MS. Prognostic value of pre-transplant mean pulmonary arterial pressure in lung transplant recipients: a single-institution experience. *J Thorac Dis* 2018;10(3):1578-1587. doi: 10.21037/jtd.2018.03.46